

Subelliptic estimates for some systems of complex vector fields : quasihomogeneous case

M. Derridj

5 rue de la Juvinière

78 350 Les loges en Josas, France.

and

B. Helffer

Laboratoire de Mathématiques, Univ Paris-Sud and CNRS,

F 91 405 Orsay Cedex, France.

February 2, 2008

Abstract

For about twenty five years it was a kind of folk theorem that complex vector-fields defined on $\Omega \times \mathbb{R}_t$ (with Ω open set in $\mathbb{R}^n$) by

$$L_j = \frac{\partial}{\partial t_j} + i \frac{\partial \varphi}{\partial t_j}(\mathbf{t}) \frac{\partial}{\partial x}, \quad j = 1, \dots, n, \quad \mathbf{t} \in \Omega, x \in \mathbb{R},$$

with φ analytic, were subelliptic as soon as they were hypoelliptic. This was the case when $n = 1$ but in the case $n > 1$, an inaccurate reading of the proof given by Maire (see also Trèves) of the hypoellipticity of such systems, under the condition that φ does not admit any local maximum or minimum (through a non standard subelliptic estimate), was supporting the belief for this folk theorem. Quite recently, J.L. Journé and J.M.Trépreau show by examples that there are very simple systems (with polynomial φ 's) which were hypoelliptic but not subelliptic in the standard L^2 -sense. So it is natural to analyze this problem of subellipticity which is in some sense intermediate (at least when φ is C^∞) between the maximal hypoellipticity (which was analyzed by Helffer-Nourrigat and Nourrigat) and the simple local hypoellipticity (or local microhypoellipticity) and to start first with the easiest non trivial examples. The analysis presented here is a continuation of a previous work by the first author and is devoted to the case of quasihomogeneous functions.

1 Introduction and Main result

1.1 Preliminaries

Let Ω an open set in $\mathbb{R}^n$ with $0 \in \Omega$. We consider the regularity properties of the following family of complex vector fields on $\Omega \times \mathbb{R}$

$$L_j = \frac{\partial}{\partial t_j} + i \frac{\partial \varphi}{\partial t_j}(\mathbf{t}) \frac{\partial}{\partial x}, \quad j = 1, \dots, n, \quad \mathbf{t} \in \Omega, x \in \mathbb{R}, \quad (1.1)$$

where $\varphi \in C^1(\Omega, \mathbb{R})$, with $\varphi(0) = 0$. We will concentrate our analysis near a point $(0, 0)$ (but note that the operator is invariant by translation in the x variable).

Many authors have considered this type of systems. For a given Ω , they were in particular interested in the existence, for some pair (s, N) such that $s + N > 0$, of the following family of inequalities.

For any pair of bounded open sets ω, I such that $\overline{\omega} \subset \Omega$ and $I \subset \mathbb{R}$, there exists a constant $C_{s,N}(\omega, I)$ such that

$$\|u\|_s^2 \leq C_N(\omega, I) \left(\sum_{j=1}^n \|L_j u\|_0^2 + \|u\|_{-N}^2 \right), \quad \forall u \in C_0^\infty(\omega \times I), \quad (1.2)$$

where $\|\cdot\|_r$ denotes the Sobolev norm in $H^r(\Omega \times \mathbb{R})$.

If $s > 0$, we say that we have a subelliptic estimate. In [JoTre], there are also results where s can be arbitrarily negative. We will speak in the case when $s \leq 0$ of weak subellipticity. Note that in this case the existence of this inequality is not sufficient for proving hypoellipticity.

The system (1.1) being elliptic in the $\mathbf{t}$ variable, it is enough to analyze the subellipticity microlocally near $\tau = 0$, i.e. near $(0, (0, \xi))$ in $(\omega \times I) \times (\mathbb{R}^{n+1} \setminus \{0\})$ with $\{\xi > 0\}$ or $\{\xi < 0\}$.

This leads to the analysis of the existence of two constants C_s^+ and C_s^- such that the two following inequalities hold :

$$\int_{\omega \times \mathbb{R}^+} \xi^{2s} |\widehat{u}(\mathbf{t}, \xi)|^2 d\mathbf{t} d\xi \leq C_s^+ \int_{\omega \times \mathbb{R}^+} |\widehat{Lu}(\mathbf{t}, \xi)|^2 d\mathbf{t} d\xi, \quad \forall u \in C_0^\infty(\omega \times \mathbb{R}), \quad (1.3)$$

where $\widehat{u}(\mathbf{t}, \xi)$ is the partial Fourier transform of u with respect to the x variable, and

$$\int_{\omega \times \mathbb{R}^-} |\xi|^{2s} |\widehat{u}(\mathbf{t}, \xi)|^2 d\mathbf{t} d\xi \leq C_s^- \int_{\omega \times \mathbb{R}^-} |\widehat{Lu}(\mathbf{t}, \xi)|^2 d\mathbf{t} d\xi, \quad \forall u \in C_0^\infty(\omega \times \mathbb{R}). \quad (1.4)$$

When (1.3) is satisfied, we will speak of microlocal subellipticity in $\{\xi > 0\}$ and similarly when (1.4) is satisfied, we will speak of microlocal subellipticity in $\{\xi < 0\}$. Of course, when $s > 0$, it is standard that these two inequalities imply (1.2).

We now observe that (1.3) for φ is equivalent to (1.4) for $-\varphi$, so it is enough to concentrate our analysis on the first case.

1.2 The main result

In [De], the first author gave a sufficient condition on φ for getting (1.2) with $s > 0$. In this article, we consider the case of quasihomogeneous functions φ on $\mathbb{R}^2$ (i.e. $n = 2$) and we will give a simple condition of subellipticity where s will be related rather simply with the quasihomogeneity of φ .

These conditions will be expressed for φ in C^1 but note that they become more simple in the analytic case (see Section 7).

More precisely, let m and ℓ in $\mathbb{R}^+$ such that

$$\ell \geq 1 , \tag{1.5}$$

and

$$m \geq 2\ell . \tag{1.6}$$

We make these two assumptions¹ in the whole paper.

We consider in $\mathbb{R}^2$ (t, s) as the variables (instead of $\mathbf{t}$) and the functions $\varphi \in C^1(\mathbb{R}^2)$ will be quasihomogeneous in the following sense

$$\varphi(\lambda t, \lambda^\ell s) = \lambda^m \varphi(t, s) , \quad \forall (t, s, \lambda) \in \mathbb{R}^2 \times \mathbb{R}^+ . \tag{1.7}$$

According to (1.7), the real function φ is determined by its restriction $\tilde{\varphi}$ to the distorted circle $\mathcal{S}$

$$\tilde{\varphi} := \varphi|_{\mathcal{S}} . \tag{1.8}$$

where $\mathcal{S}$ is defined by

$$\mathcal{S} = \{(t, s) ; t^{2\ell} + s^2 = 1\} , \tag{1.9}$$

Our main result is stated under the following assumption on $\tilde{\varphi}$.

¹ with in addition the assumption that ℓ is rational in the analytic case,

Assumption 1.1 (H2)

- (i) $\tilde{\varphi}$ is not strictly negative.
- (ii) $\tilde{\varphi}$ cannot have, at any of its zeroes, a local maximum.
- (iii) If $\mathcal{S}_j^+$ is a connected component of $\tilde{\varphi}^{(-1)}(]0, +\infty[)$, then one can write $\mathcal{S}_j^+$ as a finite union of arcs satisfying Property 1.2 below.
- (iv) If $\mathcal{S}_j^-$ is one connected component of $\tilde{\varphi}^{(-1)}(]-\infty, 0])$, then $\tilde{\varphi}$ has a unique minimum in $\mathcal{S}_j^-$.
- (v) There exists $p \geq 1$, such that, if θ_0 is a zero of $\tilde{\varphi}$, then there exists an open arc $\mathcal{V}_{\theta_0}$ containing θ_0 in $\mathcal{S}$ and $C_0 > 0$, such that

$$|\tilde{\varphi}(\theta) - \tilde{\varphi}(\theta')| \geq \frac{1}{C_0} |\theta - \theta'|^p, \quad \forall \theta, \theta' \in \mathcal{V}_{\theta_0}, \quad (1.10)$$

with θ and θ' in the same connected component in $\mathcal{V}_{\theta_0} \setminus \{\theta_0\}$.

Here in the third item of (H2), we mean by saying that a closed arc $[\theta, \theta']$ has Property 1.2 the following :

Property 1.2

There exists on this arc a point $\hat{\theta}$ such that :

- (a) $\tilde{\varphi}$ is non decreasing on the arc $[\theta, \hat{\theta}]$ and non increasing on the arc $[\hat{\theta}, \theta']$. (So the restriction of $\tilde{\varphi}$ to $[\theta, \theta']$ has a maximum at $\hat{\theta}$).
- (b)

$$\langle \hat{\theta} | \theta \rangle_\ell \geq 0 \text{ and } \langle \hat{\theta} | \theta' \rangle_\ell \geq 0, \quad (1.11)$$

where for $\theta = (\alpha, \beta)$ and $\hat{\theta} = (\hat{\alpha}, \hat{\beta})$ in $\mathcal{S} \subset \mathbb{R}^2$,

$$\langle \hat{\theta} | \theta \rangle_\ell := \hat{\alpha} \alpha |\hat{\alpha}|^{\ell-1} |\alpha|^{\ell-1} + \hat{\beta} \beta. \quad (1.12)$$

Note here that we could have $\tilde{\varphi}$ constant on $\mathcal{S}_{\theta, \theta'}$ and $\hat{\theta} = \theta$ or θ' . Moreover item (b) says that the length of the two arcs is sufficiently small, more precisely that the distorted “angles” (see Section 3) associated to $[\theta, \theta']$ are acute.

We can now state our main theorem :

Theorem 1.3

Let $\varphi \in C^1(\mathbb{R}^2, \mathbb{R})$ satisfying (1.7), with ℓ and m satisfying (1.5) and (1.6). Then L is microlocally subelliptic in $\{\xi > 0\}$.

Remarks 1.4

- (i) The proof of Theorem 1.3 consists in showing that Assumption (H2) implies the assumption $(H_+(\alpha))$ with $\alpha = \frac{1}{\max(m,p)}$ introduced in [De] and which will be recalled in Section 2.
- (ii) [De] was considering the homogeneous case $\ell = 1$ and $m \geq 2$. Here we generalize by considering the quasihomogeneous case but the sufficient condition given here for getting Assumption 2.1 satisfied will be already an improvement in the homogeneous case.
- (iii) The conditions on $\tilde{\varphi}$ are more restrictive on the arcs S_k^- .
- (iv) If φ is analytic and ℓ is rational, the statement of the main theorem becomes simpler. (iii) and (v) are indeed automatically satisfied as soon that $\tilde{\varphi}$ is not identically 0. Moreover, if we write $\ell = \frac{\ell_2}{\ell_1}$ (with ℓ_1 and ℓ_2 mutually prime integers), all the criteria involving $\tilde{\varphi}$ can be reinterpreted as criteria involving the restriction $\hat{\varphi}$ of φ on

$$\mathcal{S}_{\ell_1, \ell_2} = \{(t, s) ; t^{2\ell_2} + s^{2\ell_1} = 1\} .$$

- (v) The condition (H2), ii) is natural and cannot be relaxed according to the necessary conditions obtained by J.L. Journé and J.M. Trépreau [JoTre] for the subellipticity of these systems.

Organization of the paper

The proof of our main theorem will be based on a “abstract” criterion established in [De], which will be recalled in Section 2. After the introduction of a terminology adapted to the quasihomogeneity of the problem in Section 3, we continue with the proof of the general main theorem in three steps starting from the analysis of the quasielliptic case in Section 4, showing then how one can localize the proof in suitable quasihomogeneous sectors in Section 5 and finishing by proving the general case in Section 6. Section 7 is devoted to the particular case of an analytic function φ . We give in an appendix the computation of a basic Jacobian, whose control is important in the verification of the abstract criterion.

Acknowledgements

At various stages of this work, the authors have fruitful discussions with F. Nier, J.M. Trépreau and H. Maire.

2 Derridj's subellipticity criterion.

2.1 The statement

We now recall the criterion established in [De]. This involves, for a given $\alpha > 0$, the following geometric escape condition on φ . We do not have in this section the restriction $n = 2$

Assumption 2.1 ($H_+(\alpha)$)

There exist an open set $\omega \subset \Omega$ and $\tilde{\omega} \subset \omega$, with full Lebesgue measure in ω , and a map $\gamma :$

$$\tilde{\omega} \times [0, 1] \ni (\mathbf{t}, \tau) \mapsto \gamma(\mathbf{t}, \tau) \in \Omega ,$$

such that

$$(i) \quad \gamma(\mathbf{t}, 0) = \mathbf{t} ; \gamma(\mathbf{t}, 1) \notin \omega , \forall \mathbf{t} \in \tilde{\omega} .$$

$$(ii) \quad \gamma \text{ is of class } C^1 \text{ outside a negligible set } E \text{ and there exist } C_1 > 0, \\ C_2 > 0 \text{ and } C_3 > 0 \text{ such that}$$

$$(a)$$

$$|\partial_\tau \gamma(\mathbf{t}, \tau)| \leq C_2 , \forall (\mathbf{t}, \tau) \in \tilde{\omega} \times [0, 1] \setminus E .$$

$$(b)$$

$$|\det(D_{\mathbf{t}}\gamma)(\mathbf{t}, \tau)| \geq \frac{1}{C_1} , \tag{2.1}$$

where $D_{\mathbf{t}}\gamma$ denotes the Jacobian matrix of γ considered as a map from $\tilde{\omega}$ into $\mathbb{R}^2$.

$$(c)$$

$$\varphi(\gamma(\mathbf{t}, \tau)) - \varphi(\mathbf{t}) \geq \frac{1}{C_3} \tau^\alpha , \forall (\mathbf{t}, \tau) \in \tilde{\omega} \times [0, 1] .$$

Using this assumption, it is proved in [De] the following theorem.

Theorem 2.2

If φ satisfies ($H_+(\alpha)$), then the associated system $(1.1)_\varphi$ is microlocally

$\frac{1}{\alpha}$ -subelliptic in $\{\xi > 0\}$.

If $-\varphi$ satisfies $(H_+(\alpha))$, then the associated system $(1.1)_\varphi$ is microlocally

$\frac{1}{\alpha}$ -subelliptic in $\{\xi < 0\}$.

If φ and $-\varphi$ satisfy $(H_+(\alpha))$, then the associated system $(1.1)_\varphi$ is $\frac{1}{\alpha}$ -subelliptic.

2.2 The proof

For the commodity of the reader, we reproduce the proof of [De] in the case $\xi > 0$.

If $u \in C_0^\infty(\omega \times \mathbb{R})$, one can, using the partial Fourier transform (with respect to x), recover u from $f = Lu$ by

$$\widehat{u}(\mathbf{t}, \xi) = - \int_0^1 \exp [\xi \varphi(\gamma(\mathbf{t}, \tau))] \widehat{f}(\gamma(\mathbf{t}, \tau), \xi) \cdot (\partial_\tau \gamma)(\mathbf{t}, \tau) d\tau . \quad (2.2)$$

Taking Cauchy-Schwarz in (2.2), we obtain

$$\begin{aligned} |\widehat{u}(\mathbf{t}, \xi)|^2 &\leq \\ &\left(\int_0^1 \exp [\xi \varphi(\gamma(\mathbf{t}, \tau))] d\tau \right) \times \\ &\quad \times \left(\int_0^1 \exp [\xi \varphi(\gamma(\mathbf{t}, \tau))] |\widehat{f}(\gamma(\mathbf{t}, \tau), \xi) \cdot (\partial_\tau \gamma)(\mathbf{t}, \tau)|^2 d\tau \right) . \end{aligned} \quad (2.3)$$

By items $(ii)_{(b)}$ and $(ii)_{(c)}$ in Assumption 2.1, this implies

$$|\widehat{u}(\mathbf{t}, \xi)|^2 \leq C_2^2 \left(\int_0^1 \exp -\frac{1}{C_3} \tau^\alpha \xi d\tau \right) \times \left(\int_0^1 \exp -\frac{1}{C_3} \tau^\alpha \xi |\widehat{f}(\gamma(\mathbf{t}, \tau), \xi)|^2 d\tau \right) .$$

So, integrating in $\mathbf{t}$ over ω , we get

$$\begin{aligned} &\int_\omega |\widehat{u}(\mathbf{t}, \xi)|^2 d\mathbf{t} \\ &\leq C_2^2 \left(\int_0^1 \exp -\frac{1}{C_3} \tau^\alpha \xi d\tau \right) \times \left(\int_0^1 \int_\omega \exp -\frac{1}{C_3} \tau^\alpha \xi |\widehat{f}(\gamma(\mathbf{t}, \tau), \xi)|^2 d\mathbf{t} d\tau \right) . \end{aligned} \quad (2.4)$$

We now change of variables : $\mathbf{v} = \gamma(\mathbf{t}, \tau)$, $\tau = \tau$. The second term in the r.h.s. of (2.4) can be estimated as follows.

$$\begin{aligned} &\int_0^1 \int_\omega (\exp -\frac{1}{C_3} \tau^\alpha \xi) |\widehat{f}(\gamma(\mathbf{t}, \tau), \xi)|^2 d\mathbf{t} d\tau \\ &= \int_0^1 \int_\omega (\exp -\frac{1}{C_3} \tau^\alpha \xi) |\widehat{f}(\mathbf{v}, \xi)|^2 |D_{\mathbf{t}} \gamma|^{-1} d\tau d\mathbf{v} \\ &\leq C_1 \left(\int_0^1 \exp -\frac{1}{C_3} \tau^\alpha \xi d\tau \right) \left(\int_\Omega |\widehat{f}(\mathbf{v}, \xi)|^2 d\mathbf{v} \right) , \end{aligned}$$

where we have used the lower bound (2.1) for the Jacobian $|D_{\mathbf{t}} \gamma|$.

So finally, one has

$$\int_{\omega} |\widehat{u}(\mathbf{t}, \xi)|^2 d\mathbf{t} \leq C_1 C_2^2 \left(\int_0^1 \left(\exp - \frac{1}{C_3} \tau^\alpha \xi \right) d\tau \right)^2 \left(\int_{\Omega} |\widehat{f}(\mathbf{v}, \xi)|^2 d\mathbf{v} \right).$$

We then obtain the existence of $C(C_1, C_2, C_3) > 0$ such that, for all $\xi > 0$,

$$\int_{\omega} |\widehat{u}(\mathbf{t}, \xi)|^2 d\mathbf{t} \leq C(C_1, C_2, C_3) |\xi|^{-\frac{2}{\alpha}} \left(\int_{\Omega} |\widehat{f}(\mathbf{v}, \xi)|^2 d\mathbf{v} \right).$$

3 Quasihomogeneous structure

3.1 Distorted geometry

Condition (i) in Assumption 2.1 expresses the property that the curve is escaping from ω . For the description of escaping curves, it appears useful to extend the usual terminology used in the Euclidean space $\mathbb{R}^2$ in a way which is adapted to the given quasihomogeneous structure. This is realized by introducing the *dressing* map :

$$(t, s) \mapsto d_{\ell}(t, s) = (t |t|^{\ell-1}, s) . \quad (3.1)$$

which is at least of class C^1 as $\ell \geq 1$, and whose main role is to transport the distorted geometry onto the Euclidean geometry.

The first example was the unit distorted circle (in short unit disto-circle or unit “circle”) $\mathcal{S}$ introduced in (1.9) whose image by d_{ℓ} becomes the standard unit circle in $\mathbb{R}^2$ centered at $(0, 0)$.

Similarly, we will speak of disto-sectors, disto-arcs, disto-rays, disto-disks. In particular, for $(a, b) \in \mathcal{S}$, we define the disto-ray $\mathcal{R}_{(a,b)}$ by

$$\mathcal{R}_{(a,b)} := \{(\lambda a, \lambda^{\ell} b) ; 0 \leq \lambda \leq 1\} . \quad (3.2)$$

The disto-scalar product of two vectors in $\mathbb{R}^2$ (t, s) et (t', s') is then given by

$$\langle (t, s) | (t', s') \rangle_{\ell} = tt' |tt'|^{\ell-1} + ss' . \quad (3.3)$$

(for $\ell = 1$, we recover the standard scalar product).

For $(t, s) \in \mathbb{R}^2$, we introduce also the quasihomogeneous positive function ϱ defined on $\mathbb{R}^2$ by :

$$\varrho(t, s)^{2\ell} = t^{2\ell} + s^2 . \quad (3.4)$$

With these notations, we observe that, if $(t, s) \in \mathbb{R}^2 \setminus \{(0, 0)\}$, then

$$(\tilde{t}, \tilde{s}) := \left(\frac{t}{\varrho(t, s)}, \frac{s}{\varrho(t, s)} \right) \in \mathcal{S}, \quad (3.5)$$

and

$$(t, s) \in \mathcal{R}_{(\tilde{t}, \tilde{s})}.$$

The open disto-disk $D(R)$ is then defined by

$$D(R) = \{(x, y) \mid \varrho(x, y) < R\}. \quad (3.6)$$

We can also consider a parametrization of the disto-circle by a parameter on the corresponding circle $\vartheta \in \mathbb{R}/2\pi\mathbb{Z}$ (through the dressing map). We note that we have a natural (anticlockwise) orientation of the disto-circle. In other cases it will be better to parametrize by s (if $t \neq 0$) or by t (if $s \neq 0$). So a point in $\mathcal{S}$ will be defined either by θ or by $(a, b) \in \mathbb{R}^2$ or by ϑ .

Once an orientation is defined on $\mathcal{S}$, two points θ_1 and θ_2 (or (a_1, b_1) and (a_2, b_2)) on $\mathcal{S}$ will determine a unique unit “sector” $V \subset D(1)$.

3.2 Distorted dynamics

The parametrized curves γ permitting us to satisfy Assumption 2.1 will actually be “lines” (possibly broken) which will finally escape from a neighborhood of the origin. Our aim in this subsection is to define these “lines” (actually distorted parametrized lines).

In parametric coordinates, with

$$t(\tau) = t + \varrho\tau, \quad (3.7)$$

the curve γ starting from (t, s) and disto-parallel to (c, d) is defined by writing that the vector $(t(\tau)|t(\tau)|^{\ell-1} - t|t|^{\ell-1}, s(\tau) - s)$ is parallel to $(c|c|^{\ell-1}, d)$.

In the applications, we will consider only consider $\varrho = \pm c$.

So

$$(t(\tau)|t(\tau)|^{\ell-1} - t|t|^{\ell-1})d = c|c|^{\ell-1}(s(\tau) - s),$$

and we find

$$s(\tau) = s + \frac{d}{c|c|^{\ell-1}} (t(\tau)|t(\tau)|^{\ell-1} - t|t|^{\ell-1}), \quad (3.8)$$

We consider the map $\sigma \mapsto f_\ell(\sigma)$ which is defined by

$$f_\ell(\sigma) = \sigma |\sigma|^{\ell-1} . \quad (3.9)$$

Note that

$$f'_\ell(\sigma) = \ell |\sigma|^{\ell-1} \geq 0 . \quad (3.10)$$

With this new function, (3.8) can be written as

$$df_\ell(t(\tau)) - s(\tau)f_\ell(c) = df_\ell(t) - sf_\ell(c) . \quad (3.11)$$

This leads us to use the notion of distorted determinant of two vectors in $\mathbb{R}^2$. For two vectors $v := (v_1, v_2)$ and $w := (w_1, w_2)$, we introduce :

$$\Delta_\ell(v; w) = f_\ell(v_1)w_2 - v_2f_\ell(w_1) . \quad (3.12)$$

We will also write :

$$\Delta_\ell(v; w) = \Delta_\ell(v_1, v_2, w_1, w_2) . \quad (3.13)$$

With these notations, (3.11) can be written

$$\Delta_\ell((c, d); (t(\tau), s(\tau))) = \Delta_\ell((c, d); (t, s)) , \quad (3.14)$$

or

$$\Delta_\ell(c, d, t(\tau), s(\tau)) = \Delta_\ell(c, d, t, s) , \quad (3.15)$$

When $\ell = 1$, we recover the usual determinant of two vectors in $\mathbb{R}^2$. For $\ell \geq 1$, we have simply the relation :

$$\Delta_\ell(v; w) = \Delta_1(d_\ell(v); d_\ell(w)) . \quad (3.16)$$

Note that $\Delta_\ell(v; w)$ vanishes when $d_\ell(v)$ and $d_\ell(w)$ are collinear.

We now look at the variation of ψ which is defined (for a given initial point (t, s)) by

$$\tau \mapsto \psi(\tau) = \rho(\tau)^{2\ell} = t(\tau)^{2\ell} + s(\tau)^2 . \quad (3.17)$$

Easy computations give also :

$$\psi'(\tau) = \frac{2\varrho}{f_\ell(c)} f'_\ell(t + \varrho\tau) \langle (c, d) \mid (t(\tau), s(\tau)) \rangle_\ell , \quad (3.18)$$

whose sign is the product of the sign of the (disto)-scalar product of (c, d) and $(t(\tau), s(\tau))$ and the sign of $c\varrho$.

We now analyze the variation of the (disto)-scalar product $\langle (c, d) \mid (t(\tau), s(\tau)) \rangle_\ell$ as a function of τ . We have the formula

$$\langle (c, d) \mid (t(\tau), s(\tau)) \rangle_\ell = \langle (c, d) \mid (t, s) \rangle_\ell + \frac{1}{f_\ell(c)} (f_\ell(t(\tau)) - f_\ell(t)) . \quad (3.19)$$

If we now assume that

$$c\varrho > 0 , \quad \langle (c, d) \mid (a, b) \rangle_\ell \geq 0 , \quad (3.20)$$

Then for (s, t) in the unit sector $\mathcal{V}_{abcd}$ associated to the arc $((a, b), (c, d))$, we obtain :

$$\psi'(\tau) \geq \frac{1}{f_\ell(c)^2} \times (2\varrho f'_\ell(t + \varrho\tau)) \times (f_\ell(t(\tau)) - f_\ell(t)) .$$

We rewrite this inequality in the form

$$\psi'(\sigma) \geq \frac{1}{f_\ell(c)^2} \times ((f_\ell(t(\sigma)) - f_\ell(t))^2)' , \quad \forall \sigma \geq 0 .$$

Integrating over $[0, \tau]$, we get for $\tau \geq 0$:

$$\psi(\tau) \geq \frac{1}{f_\ell(c)^2} \times (f_\ell(t(\tau)) - f_\ell(t))^2 . \quad (3.21)$$

We now need the following

Lemma 3.1

For any $\ell \geq 1$, $\tau \geq 0$, and $\gamma \in \mathbb{R}$, we have

$$f_\ell(\tau + \gamma) - f_\ell(\gamma) \geq f_\ell\left(\frac{\tau}{2}\right) . \quad (3.22)$$

Remark 3.2

This lemma can be improved when $\gamma \geq 0$; we can then show

$$f_\ell(\tau + \gamma) - f_\ell(\gamma) \geq f_\ell(\tau) . \quad (3.23)$$

Proof

By the previous remark, the proof is clear when $\gamma \geq 0$ or when $\gamma + \tau \leq 0$. So it remains to analyze the case when $-\tau < \gamma < 0$. But the two terms on the left hand side are now positive. So we immediately obtain (3.22) by observing that $\max(\tau + \gamma, -\gamma) \geq \frac{\tau}{2}$.

Remark 3.3

If we take the square, we obtain (and this time for any $\tau \in \mathbb{R}$) the inequality

$$(f_\ell(\tau + \gamma) - f_\ell(\gamma))^2 \geq \left(\frac{\tau}{2}\right)^{2\ell}. \quad (3.24)$$

Now using Lemma 3.1, this leads to

Lemma 3.4

Under Condition (3.20), we have, for any $\tau \geq 0$, for any $(t, s) \in \mathcal{V}_{abcd}$,

$$\rho(\tau)^{2\ell} - \rho(0)^{2\ell} \geq \left(\frac{\varrho\tau}{2c}\right)^{2\ell}. \quad (3.25)$$

If instead $\varrho c < 0$, we obtain :

$$\rho(\tau)^{2\ell} - \rho(0)^{2\ell} \leq -\left(\frac{\varrho\tau}{2c}\right)^{2\ell}. \quad (3.26)$$

We continue by analyzing the variation of $s(\tau)$ and $t(\tau)$ and more precisely the variation on the disto-circle of :

$$\tilde{t}(\tau) = \frac{t(\tau)}{\rho(\tau)}, \quad \tilde{s}(\tau) = \frac{s(\tau)}{\rho(\tau)^\ell}. \quad (3.27)$$

After some computations, we get, with

$$\varrho = \pm c,$$

$$\tilde{t}'(\tau) = \pm |c|^{1-\ell} \frac{s(\tau)}{\rho(\tau)^{2\ell+1}} \Delta_\ell(c, d, t, s), \quad (3.28)$$

which can also be written in the form

$$\tilde{t}'(\tau) = \pm |c|^{1-\ell} \frac{\tilde{s}(\tau)}{\rho(\tau)} \Delta_\ell(c, d, \tilde{t}(\tau), \tilde{s}(\tau)). \quad (3.29)$$

Similarly, we get for $\tilde{s}'$,

$$\tilde{s}'(\tau) = \mp \ell |c|^{1-\ell} \frac{t(\tau)^{2\ell-1}}{\rho(\tau)^{3\ell}} \Delta_\ell(c, d, t, s), \quad (3.30)$$

and

$$\tilde{s}'(\tau) = \mp \ell |c|^{1-\ell} \frac{\tilde{t}^{2\ell-1}(\tau)}{\rho(\tau)} \Delta_\ell(c, d, \tilde{t}(\tau), \tilde{s}(\tau)). \quad (3.31)$$

4 Analysis of the quasielliptic case ($\tilde{\varphi} > 0$)

We first start the proof of the main theorem with the particular case when

$$\tilde{\varphi} \geq \mu > 0. \quad (4.1)$$

This case is already interesting for presenting the main ingredients of the general proof. We can remark indeed that what we are doing below in $\mathcal{S}$ can be done later in a specific (disto)-arc of $\mathcal{S}$.

On $\mathcal{S}$, taking a regular parametrization of $\mathcal{S}$ denoted by θ , we consider the connected components of $\{\tilde{\varphi}' \geq 0\}$ or of $\{\tilde{\varphi}' \leq 0\}$.

Our assumption takes in this case the following form.

Assumption 4.1 (H_1)

Either the cardinal of the connected components in $\mathcal{S}$ of $\{\tilde{\varphi}' \geq 0\}$ or the cardinal of the connected components of $\{\tilde{\varphi}' \leq 0\}$ is finite.

Remark 4.2

This assumption is automatically satisfied if $\tilde{\varphi}$ is analytic.

Theorem 4.3

Let φ in $C^1(\mathbb{R}^2; \mathbb{R})$ satisfying (1.7) and (4.1). Then (H_1) implies $(H_+(\alpha))$, with $\alpha = \frac{1}{m}$.

Corollary 4.4

If (H_1) is satisfied, then the system (1.1) is $\frac{1}{m}$ -subelliptic in $\{\xi > 0\}$.

Proof of Theorem 4.3

Step 1 : Construction of the covering

Under Assumption (H_1) (say for definiteness under the first alternative), and considering the connected components introduced there, we first start by constructing a finite covering of $\mathcal{S}$ by a family of disjoint open arcs $\mathcal{S}_j$ such that

$$\mathcal{S} = \cup_j \overline{\mathcal{S}_j}, \quad (4.2)$$

in the following way. If θ_j (or (a_j, b_j)) denotes the sequence of the left end points of the components of $\{\tilde{\varphi}' \geq 0\}$ and by $\hat{\theta}_j$ (or (c_j, d_j)) the sequence of the right end points, we define $\mathcal{S}_j$ as the arc $\mathcal{S}_j = (\theta_j, \theta_{j+1})$. We observe

that :

$\tilde{\varphi}$ is non decreasing on $(\theta_j, \hat{\theta}_j)$ and strictly decreasing on $(\hat{\theta}_j, \theta_{j+1})$.

We now associate to the (disto)-arc $\mathcal{S}_j$ its (disto)-unit sector $\mathcal{V}_j$. For technical reasons, we will add a finite number of points such that we finally a (possibly new) family of open arcs $\mathcal{S}_j = (\theta_j, \theta_{j+1})$ such that each arc satisfies Property 1.2.

Step 2 : Construction of γ

We construct γ independently in each sector $\mathcal{V}_j$. More precisely, this will be a map from $[(\mathcal{V}_j \cup \mathcal{R}_{\theta_j}) \setminus \{(0, 0)\}] \times [0, 1]$ into Ω (and actually in the infinite sector associated to $\mathcal{S}_j$). From now on in this paragraph, we fixed j (and take it equal to 1). So $\mathcal{S}_1$ denotes the set of points $(t, s) \in D(1)$ such that $(\frac{t}{\rho(t, s)}, \frac{s}{\rho(t, s)^\ell}) \in (\theta_1, \theta_2)$.

We now define γ (see (3.7)-(3.8), with $\varrho = c_1$) by

(i) If $c_1 \neq 0$,

$$\gamma(t, s, \tau) := (t(\tau), s(\tau)) := (t + c_1\tau, s + \frac{d_1}{f_\ell(c_1)}(f_\ell(t(\tau)) - f_\ell(t))) . \quad (4.3)$$

(ii) If $c_1 = 0$

$$\gamma(t, s, \tau) := (t(\tau), s(\tau)) := (t, s + d_1\tau) . \quad (4.4)$$

Remarks 4.5

(i) Note that for any (t_0, s_0, τ) the Jacobian of the map $(t, s) \mapsto \gamma(t, s, \tau)$ at (t_0, s_0) is 1.

(ii) Actually, one can avoid the second case by replacing $c_1 = 0$ by an arbitrarily close $\tilde{c}_1$ whose sign will depend on the considered “sector”.

From now on, we write for simplification $c = c_1, d = d_1$.

Let us look at the most generic case² when $c \neq 0$. In order to show $(H_+(\alpha))$, the only non trivial property is to show property (ii) (b) in Assumption 2.1.

²We will add in footnotes some indications for the other case.

Lemma 4.6

If (t, s) belongs to the “subsector” associated with $[\theta_1, \widehat{\theta}_1]$ (resp. to the “subsector” associated with $[\widehat{\theta}_1, \theta_2]$), then the whole curve $\gamma(t, s, \tau)$ stays in the same (infinite) “subsector” for $\tau \geq 0$.

Proof

The lemma is geometrically evident using the dressing map d_ℓ .

Step 3 : Lower bound along the curve γ

Let us introduce a few more notations.

$$\rho(\tau) = \rho(\gamma(t, s, \tau)) , \quad \rho = \rho(0), \quad (4.5)$$

and

$$\theta(\tau) = \theta_\tau = \left(\frac{t(\tau)}{\rho(\tau)}, \frac{s(\tau)}{\rho^\ell(\tau)} \right) = (\widetilde{t}(\tau), \widetilde{s}(\tau)) , \quad (4.6)$$

with $\theta = \theta(0) = (\widetilde{t}, \widetilde{s})$.

We note that with the above notations

$$\varphi(t, s) = \rho^m \widetilde{\varphi}(\widetilde{t}, \widetilde{s}) . \quad (4.7)$$

We want to show (ii)_(c) in Assumption 2.1 and first decompose the expression which has to be estimated from below by writing :

$$\rho(\tau)^m \widetilde{\varphi}(\theta_\tau) - \rho^m \widetilde{\varphi}(\theta) = (\rho(\tau)^m - \rho^m) \widetilde{\varphi}(\theta) + \rho(\tau)^m (\widetilde{\varphi}(\theta_\tau) - \widetilde{\varphi}(\theta)) , \quad (4.8)$$

and will obtain a lower bound for each term of the r.h.s. of (4.8).

Analysis of $\rho(\tau)^m - \rho^m$.

Using (3.25) and

$$\rho(\tau)^m - \rho^m = (\rho(\tau)^{2\ell})^{\frac{m}{2\ell}} - (\rho^{2\ell})^{\frac{m}{2\ell}} \geq (\rho(\tau)^{2\ell} - \rho^{2\ell})^{\frac{m}{2\ell}} ,$$

where we note that $m \geq 2\ell$, we deduce

$$\rho(\tau)^m - \rho^m \geq 2^{-m} \tau^m , \quad \forall \tau \geq 0 . \quad (4.9)$$

Finally³, using the lower bound for $\tilde{\varphi}$, we get for the first term of the r.h.s. of (4.8) the inequality :

$$(\rho(\tau)^m - \rho^m)\tilde{\varphi}(\theta) \geq 2^{-m}\mu\tau^m . \quad (4.10)$$

Analysis of $\tilde{\varphi}(\theta_\tau) - \tilde{\varphi}(\theta)$.

Having in mind our assumptions on the variation of $\tilde{\varphi}$ on $[\theta_1, \theta_2]$, we have simply to prove here :

- If $(\tilde{t}, \tilde{s}) \in [\theta_1, \hat{\theta}_1]$, the function $\tau \mapsto \theta_\tau$ is non decreasing.
- If $(\tilde{t}, \tilde{s}) \in [\hat{\theta}_1, \theta_2]$, the function $\tau \mapsto \theta_\tau$ is non increasing.

But this is immediate after having sent the initial picture by the dressing map (3.1).

5 Control in the case of the sectors $\mathcal{V}_j^+ \cup \mathcal{R}_{(a_j, b_j)}$

Provisory assumption .

For the control, of the order of the zeroes, we assume for the moment that the p appearing in (1.10) (Assumption 1.1) satisfies

$$1 \leq p \leq m . \quad (5.1)$$

This will be removed later (see Subsection 6.3).

In comparison with the quasielliptic situation, the only point is that the condition of positivity could be not satisfied at one or two ends. Note that we keep Assumption 1.2 iii). The essential idea is then to improve the second part of the lower bound for (4.8) using the fifth item of Assumption 1.1 (i.e. the lower bound (1.10)).

Having in mind what we have done before, it remains to analyze “rays” coming from points (t, s) which are close to $\mathcal{R}_{(a_j, b_j)}$. The key estimate is that

³ In the case, when $c = 0$, then if $ds > 0$, we have

$$\psi(\tau) = (s + d\tau)^2 - s^2 \geq \tau^2 \geq \tau^{2\ell} , \quad \forall \tau \in [0, 1].$$

there exists $\mu_0 > 0$ and a “sector” $\mathcal{V}$ neighboring this “end-ray” such that

$$\tilde{\varphi}(\theta_\tau) - \tilde{\varphi}(\theta) \geq \mu_0 \left(\frac{\tau}{\rho(\tau)} \right)^m, \quad \forall (t, s) \in \mathcal{V}, \quad \forall \tau \in [0, 1]. \quad (5.2)$$

But this proof is immediate from (1.10), if we control more precisely the positivity of $\theta(\tau) - \theta(0)$, as we shall see in (5.9).

Quantitative control of the positivity

We treat three typical cases.

Case 1 : $c > 0, d > 0, a_1 > 0, b_1 < 0$

We assume that $\tilde{\varphi}(a_1, b_1) = 0$ and it is enough to control the trajectories starting from $(s, t) \in \sigma$ where the “sector” σ is defined by the condition that the corresponding $(\tilde{t}, \tilde{s})$ close to (a_1, b_1) and between (a_1, b_1) and (c, d) :

$$(t, s) \in \sigma \text{ iff } (\tilde{t}, \tilde{s}) \in (\theta_1, \theta_1 + \epsilon_1), \quad (5.3)$$

for some $\epsilon_1 > 0$.

The point here is that one has a regular parametrization of the disto-circle by the $\tilde{s}$ variable and that we stay in the half-right plan.

By Formula (3.30), we observe that for (t, s) as before, we have

$$\tilde{s}'(\tau) \geq 0, \quad \forall \tau \in [0, 1],$$

Here we have used that $\Delta_\ell(c, d, a_1, b_1) < 0$. We also get

$$\tilde{\varphi}(\theta_\tau) - \tilde{\varphi}(\theta) \geq 0. \quad (5.4)$$

We now use (3.31) and observe that $\Delta_\ell(\tilde{t}(\tau), \tilde{s}(\tau), c, d)$ is increasing as τ increases (together with $\tilde{s}(\tau)$).

Let us observe the trivial inequality

$$1 \geq \frac{t(\tau)}{\rho(\tau)} \geq c \frac{\tau}{\rho(\tau)}. \quad (5.5)$$

So there exists ρ and for any $(s, t) \in \sigma$, $\tau(t, s)$ such that

$$\Delta_\ell(\tilde{t}(\tau), \tilde{s}(\tau); (c, d)) \geq \rho,$$

for all $\tau \in [0, \tau(t, s)]$ and $\Delta_\ell(\tilde{t}(\tau), \tilde{s}(\tau); (c, d)) \leq \rho$ if $\tau \in [\tau(t, s), 1]$.
In the first interval, we observe that $\tilde{t}$ is semibounded on σ , so :

$$\tilde{s}(\tau) - \tilde{s}(0) \geq \mu_1 \frac{\tau}{\rho(\tau)}, \quad \forall \tau \in [0, \tau(t, s)] , \quad (5.6)$$

where $\mu_1 = \ell(\inf_{(t,s) \in \sigma} \tilde{t}^{2\ell-1})c^{1-\ell}\rho$.

This gives a uniform lower bound for $\tau \in [0, \tau(t, s)]$, but for $\tau \geq \tau(t, s)$ we have a uniform lower bound (if we choose ρ small enough) of $\tilde{s}(\tau) - \tilde{s}(0)$ by a constant $\alpha_\rho > 0$. This together with (5.5), gives the existence of $\mu_2 > 0$ such that

$$\tilde{s}(\tau) - \tilde{s}(0) \geq \mu_2 \frac{\tau}{\rho(\tau)}, \quad \forall \tau \in [\tau(t, s), 1] . \quad (5.7)$$

So we have finally shown that there exists $\mu_\sigma > 0$ such that

$$\tilde{s}(\tau) - \tilde{s}(0) \geq \mu_\sigma \frac{\tau}{\rho(\tau)}, \quad \forall \tau \in [0, 1] . \quad (5.8)$$

Case 2 : $c > 0, d > 0, a_2 < 0, b_2 > 0$

We assume that $\tilde{\varphi}(a_2, b_2) = 0$ and it is enough to control the trajectories starting from (s, t) with corresponding $(\tilde{s}(0), \tilde{t}(0))$ close to (a_2, b_2) and between (c, d) and (a_2, b_2) .

Here $\tilde{t}(\tau)$ may change of sign along the trajectory and we better parametrize the disto-circle by the variable $\tilde{t}$.

Here we use (3.28) for observing that $\tilde{t}(\tau)$ is this time increasing (note that $\Delta_\ell(c, d, a_2, b_2) > 0$ and $s(\tau) \geq \varrho_0 > 0$).

A similar argument to the one leading to (5.6) gives the uniform control of $|\tilde{t}(\tau) - \tilde{t}(0)|$ from below. Here we can no more use (5.5) but will instead use (3.25) which implies

$$\frac{\tau}{\rho(\tau)} \leq 2 ,$$

without to assume the positivity of $\tilde{t}$.

There are limiting cases when we shall use both parametrizations but this does not create considerable troubles. Typically, let us consider the following case.

Case 3 : $c = 0, d = -1, a_1 < 0, b_1 < 0$

We have just to use in this case vertical escaping “rays”.

So in all the cases we obtain that, if we start (inside $\mathcal{V}_j^+$) of a “subsector” σ , whose closure does not meet the “ray” $\mathcal{R}_{(c,d)}$ then, there exists $\mu_\sigma > 0$ such that

$$|\theta(\tau) - \theta(0)| \geq \mu_\sigma \frac{\tau}{\rho(\tau)}, \quad \forall \tau \in [0, 1]. \quad (5.9)$$

So we have finally proved

Lemma 5.1

Assuming that $\tilde{\varphi}$ satisfies in $\mathcal{V}_j^+$ Property 1.2 and the non degeneracy Assumption (1.10) for its zero possibly appearing at (a_j, b_j) , then there exists $\mu > 0$ such that

$$\rho(\tau)^m \varphi(\theta(\tau)) - \rho(0)^m \varphi(\theta(0)) \geq \mu \tau^m, \quad \forall \tau \in [0, 1], \quad (5.10)$$

with $(\rho(0), \theta(0))$ corresponding to a point of $\mathcal{V}_j^+ \cup \mathcal{R}_{(a_j, b_j)}$.

Remark 5.2

A similar result can be obtained for $\mathcal{V}_j^+ \cup \mathcal{R}_{(a_{j+1}, b_{j+1})}$ with a zero at (a_{j+1}, b_{j+1}) .

6 Control for the “sectors” where $\tilde{\varphi}$ is non positive

6.1 Main case

We consider first the case of small opening “sectors” $\mathcal{V}_k^-$. More precisely, we assume that

$$\langle (c, d) | (a_j, b_j) \rangle_\ell > 0, \quad \text{for } j = k, k+1. \quad (6.1)$$

6.1.1 The subcase $c \neq 0$

We start with the case when

$$c \neq 0, \quad (6.2)$$

We keep the same notations (change the labelling by taking $k = 1$) but this time (c, d) is a point of $\mathcal{S}$ where $\tilde{\varphi}$ is a minimum. The map γ_- is defined as follows.

If $(t, s) \in D(\frac{1}{4})$ is such that $(\tilde{t}, \tilde{s}) \in ((a_1, b_1), (c, d))$, we let

(i) $\gamma_- = \gamma_1$ with

$$\begin{aligned}\gamma_1(t, s, \tau) &:= (t(\tau), s(\tau)) \\ &:= \left(t - c\tau, s + \frac{d}{f_\ell(c)}(f_\ell(t(\tau)) - f_\ell(t)) \right), \quad \forall \tau \in [0, \tau_1(t, s)],\end{aligned}\tag{6.3}$$

where $\tau_1(t, s) = \tau_1$ is the smallest τ such that $\gamma_1(t, s, \tau) \in \mathcal{R}_{(a_1, b_1)}$.

(ii) $\gamma_- = \gamma_2$ with

$$\gamma_2(t, s, \tau) = \gamma_+(t(\tau_1), s(\tau_1), \tau - \tau_1), \quad \forall \tau \in [\tau_1, 1], \tag{6.4}$$

where γ_+ is the map constructed in (4.3) in the “sector” $\mathcal{V}_0^+$ (the “sector” preceding $\mathcal{V}_1$ when turning anticlockwise).




Figure 1: The escaping ray (after application of the dressing map)

If $(\tilde{t}, \tilde{s}) \in ((c, d), (a_2, b_2))$, we do the same construction with $\mathcal{R}_{(a_1, b_1)}$ replaced by $\mathcal{R}_{(a_2, b_2)}$.

Let us compute $\tau_1(t, s)$. This τ_1 is actually determined by writing that, at the corresponding point $(t(\tau_1), s(\tau_1)) = (t_1(t, s), s_1(t, s))$, we should have

$$\Delta_\ell(a_1, b_1, t_1(t, s), s_1(t, s)) = 0, \tag{6.5}$$

and

$$\Delta_\ell(c, d, t_1(t, s), s_1(t, s)) = \Delta_\ell(c, d, t, s). \tag{6.6}$$

The first one (6.5) expresses that we cross $\mathcal{R}_{a_1, b_1}$ and the second one (6.6) was observed in (3.11). This leads to the determination of τ_1 by the formula

$$f_\ell(t - c\tau_1) = f_\ell(a_1) \frac{\Delta_\ell(t, s, c, d)}{\Delta_\ell(a_1, b_1, c, d)} . \quad (6.7)$$

The uniqueness is obtained by the monotonicity of $\tau \mapsto f_\ell(t - c\tau)$ for $c \neq 0$ and the existence is a consequence of the transversality of $\mathcal{R}_{a_1, b_1}$ and the disto-parallel to $\mathcal{R}_{c, d}$ which is expressed by the condition

$$\Delta_\ell(a_1, b_1, c, d) \neq 0 . \quad (6.8)$$

It remains to control the Jacobian of the map γ .

It is immediate to see that the Jacobian of $(t, s) \mapsto \gamma(t, s, \tau)$ is equal to one when $\tau < \tau_1(t, s)$. Let us look at the more difficult case when

$$\tau > \tau_1(t, s) .$$

Under this condition, we write

$$\gamma_2(t, s, \tau) = (t_2(t, s, \tau), s_2(t, s, \tau)) ,$$

but omit the reference to τ in the computations below and prefer to think of t_1, t_2 and τ_1 as functions of (t, s) .

In order to compute the Jacobian, it is enough to compare the two 2-forms $dt_2 \wedge ds_2$ and $dt \wedge ds$.

We have in addition to (6.5) and (6.6)

$$\begin{aligned} t_1(t, s) &= t - c\tau_1(t, s) , \\ t_2(t, s) &= t_1(t, s) + \hat{c}(\tau - \tau_1(t, s)) , \\ s_2(t, s) &= s_1(t, s) + \frac{\hat{d}}{f_\ell(\hat{c})}(f_\ell(t_2(t, s)) - f_\ell(t_1(t, s))) . \end{aligned} \quad (6.9)$$

Here $(\hat{c}, \hat{d})$ denotes the coordinates (and we assume for the moment⁴ that $\hat{c} \neq 0$) of the maximum of $\tilde{\varphi}$ in the arc (corresponding to $\mathcal{V}_{k-1}^+$ in the initial notations) preceding the arc $((a_1, b_1), (a_2, b_2))$.

The two first identities imply first (by (6.5)) that

$$dt_1 \wedge ds_1 = 0 , \quad (6.10)$$

⁴See later for what we have to do when $\hat{c} = 0$.

and that (by (6.5) and (6.6)) one can express the 1-forms ds_1 and $t_1(t, s)^{\ell-1}dt_1$ as linear combinations of the 1-forms ds and $f'_\ell(t)dt$. In particular we get :

$$ds_1 = \frac{b_1}{f_\ell(a_1)} f'_\ell(t_1) dt_1 , \quad (6.11)$$

and

$$\frac{1}{b_1} \Delta_\ell(c, d, a_1, b_1) ds_1 = f_\ell(c) ds - df'_\ell(t) dt . \quad (6.12)$$

Let us start the computation of $dt_2 \wedge ds_2$ using the two last identities. We obtain (using the rules of the exterior calculus)

$$\begin{aligned} dt_2 \wedge ds_2 &= (dt_1 - \hat{c} d\tau_1) \wedge \left(ds_1 - \frac{\hat{d}}{f_\ell(\hat{c})} df_\ell(t_1(t, s)) \right) \\ &= -\hat{c} d\tau_1 \wedge \left(ds_1 - \frac{\hat{d}}{f_\ell(\hat{c})} d(f_\ell(t_1(t, s))) \right) \\ &= \delta_1 d\tau_1 \wedge ds_1 , \end{aligned}$$

with

$$\delta_1 = -|\hat{c}|^{1-\ell} b_1^{-1} \Delta_\ell(\hat{c}, \hat{d}, a_1, b_1) .$$

It remains to use the first relation of (6.9) which gives :

$$d\tau_1 = \frac{1}{c} dt - \frac{1}{c} dt_1 ,$$

and

$$dt_2 \wedge ds_2 = \frac{\delta_1}{c} dt \wedge ds_1 .$$

Using (6.12), we finally obtain the following lemma.

Lemma 6.1

If $\tau > \tau_1(t, s)$, then

$$dt_2 \wedge ds_2 = \delta dt \wedge ds , \quad (6.13)$$

with

$$\delta := -|\hat{c}|^{1-\ell} |c|^{\ell-1} \frac{\Delta_\ell(\hat{c}, \hat{d}, a_1, b_1)}{\Delta_\ell(c, d, a_1, b_1)} . \quad (6.14)$$

So the Jacobian is equal to δ , hence constant, and non zero. The fact that $\Delta_\ell(\hat{c}, \hat{d}, a_1, b_1)$ is not zero is the consequence of the assumption on the zeros of $\tilde{\varphi}$.

Remark 6.2

The existence of this lower bound of the Jacobian is probably the key point. It is shown (see [JoTre]) in the analytic case that one can always find a γ satisfying all the assumptions except this lower bound of the Jacobian by simply considering the flow associated with $\frac{1}{|\nabla\varphi|} \nabla\varphi$.

Now if we observe that $(-\varphi)$ has in $\mathcal{V}_k^-$ the properties that φ had in $\mathcal{V}_1^+$, we get rather easily the existence of $\mu > 0$ such that

$$\rho(\tau)^m \tilde{\varphi}(\theta_\tau) - \rho^m \tilde{\varphi}(\theta) \geq \mu \tau^m, \forall \tau \in [0, \tau_1(t, s)] . \quad (6.15)$$

Here we can indeed use Lemma 5.1 (after exchange of the roles of θ_τ and θ). We note also that, for $\tau > \tau_1(t, s)$, we are in a region where φ is positive so we can apply what we have done in this case. In particular, we obtain (see (4.10)) with $\tau_1 = \tau_1(t, s)$,

$$\rho(\tau)^m \tilde{\varphi}(\theta_\tau) - \rho(\tau_1) \tilde{\varphi}(\theta_{\tau_1}) \geq \mu(\tau - \tau_1)^m . \quad (6.16)$$

One can also observe (see⁵ (4.9)) that

$$\rho(\tau)^m - \rho^m \leq -\left(\frac{\tau}{2}\right)^m, \forall \tau \in [0, \tau_1(t, s)] , \quad (6.17)$$

which implies in particular the upperbound

$$\tau_1(t, s) \leq 2\rho \leq \frac{1}{2} , \quad (6.18)$$

and the inequality (see again (4.9))

$$\rho(\tau)^m - \rho(\tau_1)^m \geq \left(\frac{\tau - \tau_1}{2}\right)^m, \forall \tau \in [\tau_1, 1] . \quad (6.19)$$

From (6.18) and (6.19), one obtains that

$$\gamma(t, s, 1) \notin D\left(\frac{1}{4}\right) . \quad (6.20)$$

So the escaping condition (i) of $(H_+(\alpha))$ is satisfied with $\omega = D(\frac{1}{4})$.

On the other hand, we get from (6.15) and (6.16), the estimate

$$\rho(\tau)^m \tilde{\varphi}(\theta_\tau) - \rho^m \tilde{\varphi}(\theta) \geq 2^{-m} \mu \tau^m, \forall \tau \in [0, 1] . \quad (6.21)$$

6.1.2 The subcase $c = 0$.

For definiteness, we assume that $c = 0$ and $d = -1$ and look at initial date in the “sector” attached to the arc $((a_1, b_1), (0, -1))$. The point is just to choose a point (c', d') in $\mathcal{S}$, with $c' > 0$ and sufficiently close to $(c, d) = (0, -1)$ in order to keep the condition

$$\langle (a_1, b_1) | (c', d') \rangle > 0 .$$

We can then keep the previous construction with (c, d) replaced by (c', d') .

⁵ Here we use Assumption (6.1).

6.2 Remaining case.

In order to achieve the proof of Theorem 1.3 (i.e. to prove that $(H_+(\alpha))$ is satisfied), we have to treat the case when (6.1) is no more satisfied. So $(s, t) \mapsto \langle (c, d) | (t(\tau), s(\tau)) \rangle$ may change of sign on $\mathcal{V}_k^-$. We have avoided this problem in the case of $\mathcal{V}_k^+$ by dividing the “sector” in smaller “sectors”, but this is no more possible when $\tilde{\varphi}$ is negative. In this general case, one will be obliged to add a broken line to the two previously defined arcs (ingoing and escaping) in order to leave ω . We will see that it is always possible using a broken line made of at most five segments at the price to take ω smaller.

We start from a point (t, s) in the intersection of $\omega := D(R)$, ($R > 0$ small enough) with a “sector” associated with the arc $((a_1, b_1), (c, d))$ and we divide this “sector” into N “sectors” S^j ($j = 1, \dots, N$) of disto-angle $< \frac{\pi}{2}$. They are delimited by “rays” attached to the sequence $(\hat{\eta}_j, \hat{\zeta}_j)$ ($j = 0, \dots, N$) in $\mathcal{S}$ going **clockwise** with $(\hat{\eta}_0, \hat{\zeta}_0) = (c, d)$, $(\hat{\eta}_N, \hat{\zeta}_N) = (a_1, b_1)$.

For the commodity of the notations below, we also introduce $(\hat{\eta}_{-1}, \hat{\zeta}_{-1}) = (a_2, b_2)$ and $(\hat{\eta}_{N+1}, \hat{\zeta}_{N+1}) = (c_0, d_0)$.

It is clear that we can always do that with $N \leq 4$ and that we have previously treated the case $N = 1$.

We need also another sequence (η_j, ζ_j) in $\mathcal{S}$, which will determine the various directions of the broken line and will satisfy

$$\begin{cases} \eta_j \neq 0, \\ (\eta_j, \zeta_j) \in \left((\hat{\eta}_{j-2}, \hat{\zeta}_{j-2}), (\hat{\eta}_{j-1}, \hat{\zeta}_{j-1}) \right), \\ \Delta_\ell(\eta_j, \zeta_j, \hat{\eta}_j, \hat{\zeta}_j) \neq 0, \\ \Delta_\ell(\eta_{j+1}, \zeta_{j+1}, \hat{\eta}_j, \hat{\zeta}_j) \neq 0. \end{cases} \quad (6.22)$$

This will be satisfied by taking (η_j, ζ_j) very close (but distinct except possibly for $N = 1$ (if $c \neq 0$)) to $(\hat{\eta}_{j-1}, \hat{\zeta}_{j-1})$ for $j = 1, \dots, N$.

So we can always assure the property that starting from a point in the disk on $\mathcal{R}_{\hat{\eta}_j, \hat{\zeta}_j}$ the “straight” line parallel to $(\eta_{j+1}, \zeta_{j+1})$ will meet the “ray” $\mathcal{R}_{\hat{\eta}_{j+1}, \hat{\zeta}_{j+1}}$ inside the “disk”, for $j = 1, \dots, N - 1$.

We now explain how we construct the broken line starting from a point (t, s) belonging to the first “sector” $S_1 := S(\hat{\eta}_0, \hat{\zeta}_0, \hat{\eta}_1, \hat{\zeta}_1)$. The other cases are simpler.

Starting from (t, s) we follow for $\tau \geq 0$ the parametrized “line” parallel to (η_1, ζ_1) till we meet at the time $\tau_1(t, s)$ the “ray” $\mathcal{R}_{\hat{\eta}_1, \hat{\zeta}_1}$ at the point

$(t_1(t, s), s_1(t, s))$.

If $N > 1$ and starting now from $(t_1(t, s), t_2(t, s))$ we follow for $\tau \geq \tau_1(t, s)$ the parametrized “line” parallel to (η_2, ζ_2) till we meet at the time $\tau_2(t, s)$ the “ray” $\mathcal{R}_{\hat{\eta}_2, \hat{\zeta}_2}$ at the point $(t_2(t, s), s_2(t, s))$.

By recursion, we arrive at the point $(t_N(t, s), s_N(t, s))$ on the “ray” $R_{\hat{\eta}_N, \hat{\zeta}_N}$ at time $\tau_N(t, s)$. We can then escape along $(\eta_{N+1}, \zeta_{N+1})$.

For generalizing, what we have done for $N = 1$, we have now mainly to verify the following points :

(i) The condition of the Jacobians for $\tau \in]\tau_j(t, s), \tau_{j+1}(t, s)[$:
This will be analyzed in the appendix,

(ii) The control of the escape time :
We have to determine a condition on ω such that

$$\tau_{j+1}(t, s) - \tau_j(t, s) \leq \frac{1}{4N} , \quad (6.23)$$

(iii) The control of the dynamics :
We have to control $\rho(\tau_j(t, s))$ for $j = 1, \dots, N$, under the suitable assumption that $(s, t) \in D(0, R_N)$.

(iv) The positivity of $\tilde{\varphi}(\gamma(t, s, \tau)) - \tilde{\varphi}(t, s)$ along the trajectory :
We do not meet here new problems.

For the proof of (ii) and (iii) we observe that

$$\rho(\tau)^m + \left(\frac{\tau - \tau_i}{2} \right)^m \leq \rho(\tau_i)^m , \quad \forall \tau \in]\tau_i, \tau_{i+1}] . \quad (6.24)$$

So in particular

$$\rho(\tau_{i+1})^m + \left(\frac{\tau_{i+1} - \tau_i}{2} \right)^m \leq \rho(\tau_i)^m . \quad (6.25)$$

This gives

$$\rho(\tau_{i+1}) \leq \rho(\tau_i) , \quad (6.26)$$

and

$$\tau_{i+1} - \tau_i \leq 2\rho(\tau_i) \leq 2\rho(0) . \quad (6.27)$$

In addition, we have (see (6.19) for the case $N = 1$)

$$\rho(\tau)^m - \rho(\tau_N(t, s))^m \geq 2^{-m}(\tau - \tau_N(t, s))^m . \quad (6.28)$$

From this we deduce (without to look for optimality) that

$$R_N = \frac{1}{8} \frac{1}{N}$$

is enough for showing that the parametrized broken line has left $\omega = D(0, R_N)$ at time $\tau = 1$.

6.3 The case when $p > m$

Till now we have proved the main theorem under the additional condition (5.1). Without this additional condition, the subellipticity which is obtained is now $\inf(\frac{1}{m}, \frac{1}{p})$ instead of $\frac{1}{m}$. The only change is that we get only the existence of $\mu > 0$ such that, for $\tau \in [0, 1]$,

$$\rho(\tau)^m \varphi(\theta_\tau) - \rho^m \varphi(\theta) \geq \mu \inf \left(\tau^m, \rho(\tau)^m \left(\frac{\tau}{\rho(\tau)} \right)^p \right) . \quad (6.29)$$

But we have shown the existence of a constant $C_\varphi > 0$ and the existence of some open neighborhood of 0 ω_φ such that

$$0 \leq \tau \leq C_\varphi \rho(\tau) .$$

So we get

$$\rho(\tau)^m \varphi(\theta_\tau) - \rho^m \varphi(\theta) \geq \mu \tau^{\sup(m, p)} . \quad (6.30)$$

7 The analytic case and $\ell \in \mathbb{Q}$

7.1 The main theorem in the analytic case

We keep the previous assumptions but now assume that

$$\ell = \ell_2 / \ell_1 , \quad (7.1)$$

with ℓ_1 and ℓ_2 mutually prime integers, and that φ is analytic. In this case, the quasihomogeneity Assumption (1.7) on φ implies that φ is actually a polynomial and we can write φ in the form

$$\varphi(t, s) = \sum_{\ell_1 j + \ell_2 k = \ell_1 m} a_{j,k} t^j s^k , \quad (7.2)$$

where (j, k) are integers and the $a_{j,k}$ are real.

We can of course apply the main theorem but it is nicer to have a criterion involving more directly the assumptions on φ or on its restriction $\widehat{\varphi}$ of φ to the quasi-circle

$$\mathcal{S}_{\ell_1, \ell_2} := \{t^{2\ell_2} + s^{2\ell_1} = 1\}.$$

instead of the disto-circle $\mathcal{S}$. There are absolutely no problems if the critical points or zeroes of $\widetilde{\varphi}$ avoid $\{t = 0\} \cup \{s = 0\}$ but one should be more careful in order to analyze Condition (1.10) at the remaining points.

Let us show how this works in this case. We parametrize (this is a C^2 parametrization) on $\mathcal{S}$ by t and assume that we are close to $(0, 1)$ for definiteness and $\widetilde{\varphi}$ becomes locally with this parametrization the function

$$t \mapsto \kappa(t) = \varphi(t, \sqrt{1 - t^{2\ell}}),$$

and we assume that $\kappa(0) = 0$ and that κ is not identically 0. Suppose that we are on the side $\{t > 0\}$. Then

$$\kappa(t) = \chi(t^{\frac{1}{\ell_1}}),$$

where χ is the non identically zero analytic function

$$v \mapsto \chi(v) = \varphi(v^{\ell_1}, \sqrt{1 - v^{2\ell_2}}),$$

with $\chi(0) = 0$.

Now there exists $p > 0$ such that $\chi^{(p)}(0) \neq 0$ and we get the existence of $C > 0$ such that, in the neighborhood of 0

$$\forall v \geq 0, \forall v' \geq 0, |\chi(v) - \chi(v')| \geq \frac{1}{C}|v - v'|^p.$$

Coming back to κ , we get for a constant $\widehat{C} > 0$

$$\forall t \geq 0, \forall t' \geq 0, |\kappa(t) - \kappa(t')| \geq \frac{1}{\widehat{C}}|t^{\frac{1}{\ell_1}} - (t')^{\frac{1}{\ell_1}}|^p \geq \frac{1}{\widehat{C}}|t - t'|^p.$$

So we have obtained the proof of (1.10) for some⁶ p .

Theorem 7.1

Let φ be a real analytic non identically 0 quasihomogeneous function satisfying (1.5), (1.6), and (1.7), with $\ell = \ell_2/\ell_1$. If φ is strictly positive on $\mathbb{R}^2 \setminus \{0\}$, then φ satisfies $(H_+(\alpha))$ with $\alpha = \frac{\ell_1}{m}$ and the system (1.1) is $\frac{\ell_1}{m}$ -microlocally subelliptic in $\{\xi > 0\}$.

⁶Using that φ is a polynomial, one could get more information on p if needed.

Theorem 7.2

Let φ be a real analytic non identically 0 quasihomogeneous function satisfying (1.5), (1.6), and (1.7), with $\ell = \ell_2/\ell_1$. Suppose that φ is not a negative function. Suppose in addition that :

If $\mathcal{S}_k^- = [\theta_k, \theta_{k+1}]$ is a maximal arc where $\widehat{\varphi}$ is negative, then $\widehat{\varphi}'$ has a unique zero on $] \theta_k, \theta_{k+1}[$.

Then φ satisfies $(H_+(\alpha))$ with $\alpha > 0$. Hence the system (1.1) is microlocally subelliptic in $\{\xi > 0\}$.

Example 7.3

We recover some examples treated by H. Maire [Mai4]

$$\varphi(t, s) = t(s^2 - t^{2\ell}) , \ell \geq 1 .$$

Here $m = 2\ell + 1$, $p = 1$ and we get the subellipticity with $\alpha = \frac{1}{2\ell+1}$. As observed in [HeNi], this result is optimal and the associated system is not maximally hypoelliptic when $\ell > 1$. The maximal hypoellipticity would indeed imply $\alpha = \frac{1}{3}$.

7.2 Around Journé-Trépreau's examples

For

$$\varphi(t, s) = -t^{2m} - t^2 s^{2p} + s^q ,$$

with

$$m \geq 1 , p \geq 2 , q \geq \frac{2mp}{m-1} ,$$

J.L. Journé and J.M. Trépreau show that, although the Maire-Trèves condition is satisfied, one cannot obtain a better ρ -subellipticity than

$$\rho \leq -\left(1 - \frac{2p}{q} - \frac{1}{m}\right) \frac{n-1}{4} + \frac{1}{2q} + \frac{m-1}{4mp} .$$

The right hand side can become strictly negative. But if we impose the quasihomogeneity condition (7.2), we get as a necessary condition :

$$\left(1 - \frac{2p}{q} - \frac{1}{m}\right) = 0 ,$$

which cancels the only possible negative term.

Indeed inside this class ($m = 2$, $p = 2$), the authors can obtain the optimal subellipticity for the example

$$\varphi(t, s) = -t^4 - t^2 s^4 + s^q ,$$

with $q \geq 8$.

The optimal subellipticity is $\rho_q = \frac{3}{2q} - \frac{1}{16}$. Here let us observe that the only quasihomogeneous case corresponds to $q = 8$ and that in this case their result is coherent with our result. This example shows also that we loose the “positive” subellipticity for $q \geq 24$.

7.3 Final remarks

In the analytic case, the criterion of microhypoellipticity (proven by Maire) at say $(0, 0)$ in the direction $\{\xi > 0\}$ is that φ has no local maximum in a neighborhood of 0. When φ is quasihomogeneous, we immediately see that, at a local maximum, $\tilde{\varphi}$ should be zero. So we should avoid the following situations :

- $\tilde{\varphi} \leq 0$, with $\tilde{\varphi}^{-1}(]-\infty, 0]) \neq \emptyset$.
- $\tilde{\varphi}$ has a local maximum equal to zero on $\mathcal{S}$.

One can indeed verify that our assumptions exclude these two cases but are unfortunately more restrictive.

Note also that it would be interesting the case when $\tilde{\varphi}$ or $\hat{\varphi}$ have a strictly negative local maximum. We have not been able by our method to construct escaping curves satisfying all the conditions of the criterion in this case.

Finally, let us recall that the maximal hypoellipticity of these systems was analyzed in [HeNo, No1, No2, No3] and more recently in [HeNi].

A Control of the Jacobian for a $(N+1)$ -broken line

We just adapt the proof done for a broken line with two pieces. We treat for definiteness the case $N = 2$ and keep the same notations as in the main text.

Starting from (t, s) we consider the parametrized (disto)-line

$$t_1(\tau, t, s) = t - \eta_1 \tau, \quad s_1(\tau, t, s) = s + \frac{\zeta_1}{f_\ell(\eta_1)} (f_\ell(t_1(\tau, t, s)) - f_\ell(t)) .$$

Here (η_1, ζ_1) is a point of $\mathcal{S}$ with $\eta_1 \neq 0$ and, in our example (if in addition $c \neq 0$), $(\eta_1, \zeta_1) = (c, d)$.

We refer to (6.22) for the conditions on the sequence (η_j, ζ_j) and $(\widehat{\eta}_j, \widehat{\zeta}_j)$. We simply add the equations which are satisfied at the singular points of the broken line. The intersection with the “ray” $\mathcal{R}_{\widehat{\eta}_1, \widehat{\zeta}_1}$ occurs at time $\tau_1(t, s)$ and we get the first family of relations for the intersection point denoted by $(t_1(t, s), s_1(t, s))$.

$$\begin{aligned} \Delta_\ell(\widehat{\eta}_1, \widehat{\zeta}_1, t_1(t, s), s_1(t, s)) &= 0 ; \\ \Delta_\ell(\eta_1, \zeta_1, t_1(t, s), s_1(t, s)) &= \Delta_\ell(\eta_1, \zeta_1, t, s) ; \\ t_1(t, s) &= t - \eta_1 \tau_1(t, s) . \end{aligned} \tag{A.1}$$

Starting from $(t_1(t, s), s_1(t, s))$ we construct a new ingoing “straight”⁷ half-line $\mathcal{L}_2(t, s)$ disto-parallel to the disto-unit-vector (η_2, ζ_2) .

The straight line meets $\mathcal{R}_{\widehat{\eta}_2, \widehat{\zeta}_2}$ at time $\tau_2(t, s)$ and at the point $(t_2(t, s), s_2(t, s))$. So we get the second family of relations :

$$\begin{aligned} \Delta_\ell(\widehat{\eta}_2, \widehat{\zeta}_2, t_2(t, s), s_2(t, s)) &= 0 ; \\ \Delta_\ell(\eta_2, \zeta_2, t_2(t, s), s_2(t, s)) &= \Delta_\ell(\eta_2, \zeta_2, t_1(t, s), s_1(t, s)) ; \\ t_2(t, s) &= t_1(t, s) - \eta_2(\tau_2(t, s) - \tau_1(t, s)) . \end{aligned} \tag{A.2}$$

For $\tau > \tau_2(t, s)$, we now consider the last half-line, which is now assumed to be escaping and starting from $(t_2(t, s), s_2(t, s))$ and parallel to a new vector (η_3, ζ_3) . At time τ , we will be at the point

$$\begin{aligned} t_3(\tau, t, s) &= t_2(t, s) + \eta_3(\tau - \tau_2(t, s)) , \\ s_3(\tau, t, s) &= s_2(t, s) + \frac{\zeta_3}{f_\ell(\eta_3)} (f_\ell(t_3(\tau, t, s)) - f_\ell(t_2)) . \end{aligned} \tag{A.3}$$

⁷more precisely, straight after application of the dressing map,

It remains to control the Jacobian for the different values of $\tau \in [0, 1]$, which can be done by the computation of the coefficient of the 2-form $dt_3 \wedge ds_3$ on the 2-form $dt \wedge ds$. We will see that this coefficient is locally constant.

The proof is similar to the proof of Lemma 6.1.

We are actually able to give explicit formulas of these Jacobians, once the two sequences (η_j, ζ_j) and $(\widehat{\eta}_j, \widehat{\zeta}_j)$ are fixed.

Let us treat for definiteness the case $N = 2$ and look first at what is going on for $\tau \geq \tau_2(t, s)$.

We will show :

$$dt_3 \wedge ds_3 = \underline{\delta}_{32} d\tau_2 \wedge ds_2 = \underline{\delta}_{32} \widehat{\delta}_{21} d\tau_1 \wedge ds_1 = \delta_{3,0} dt \wedge ds . \quad (\text{A.4})$$

Let us show the existence of $\underline{\delta}_3$. Using (A.3), we first get

$$dt_3 \wedge ds_3 = dt_3 \wedge \left(ds_2 - \frac{\zeta_3}{f_\ell(\eta_3)} f'_\ell(t_2) dt_2 \right) . \quad (\text{A.5})$$

Now, the first line of (A.2), gives that

$$f_\ell(\widehat{\eta}_2) ds_2 - \widehat{\zeta}_2 f'_\ell(t_2) dt_2 = 0 . \quad (\text{A.6})$$

We then obtain easily

$$\underline{\delta}_{32} = -\eta_3 \left(1 - \frac{f_\ell(\widehat{\eta}_2) \zeta_3}{f_\ell(\eta_3) \widehat{\zeta}_2} \right) = -|\eta_3|^{1-\ell} \widehat{\zeta}_2^{-1} \Delta_\ell(\eta_3, \zeta_3, \widehat{\eta}_2, \widehat{\zeta}_2) , \quad (\text{A.7})$$

which should be non zero in our construction.

The second step is to show that

$$d\tau_2 \wedge ds_2 = \widehat{\delta}_{21} d\tau_1 \wedge ds_1 . \quad (\text{A.8})$$

The differentiation of the third line of (A.2) gives :

$$d\tau_2 \wedge ds_2 = \left(\frac{1}{\eta_2} dt_1 + d\tau_1 \right) \wedge ds_2 . \quad (\text{A.9})$$

The differentiation of the second line of (A.2) (together with (A.6)) permits us to express ds_2 as a combination of ds_1 and $f'_\ell(t_1) dt_1$.

We get indeed

$$ds_2 = \frac{\widehat{\zeta}_2}{\Delta_\ell(\eta_2, \zeta_2, \widehat{\eta}_2, \widehat{\zeta}_2)} (f_\ell(\eta_2) ds_1 - \zeta_2 f'_\ell(t_1) dt_1) ,$$

and by differentiating the first line of (A.1), we get

$$\widehat{\delta}_{21} = \frac{\widehat{\zeta}_2}{\Delta_\ell(\eta_2, \zeta_2, \widehat{\eta}_2, \widehat{\zeta}_2)} \frac{\Delta_\ell(\eta_2, \zeta_2, \widehat{\eta}_1, \widehat{\zeta}_1)}{\widehat{\zeta}_1} . \quad (\text{A.10})$$

The last step is to show

$$d\tau_1 \wedge ds_1 = \bar{\delta}_{10} dt \wedge ds .$$

The differentiation of the first line of (A.1) leads to

$$d\tau_1 \wedge ds_1 = \frac{1}{\eta_1} dt \wedge ds_1 .$$

It remains then to use the two first lines of (A.1) and we obtain

$$\bar{\delta}_{1,0} = |\eta_1|^{\ell-1} \frac{\widehat{\zeta}_1}{\Delta_\ell(\eta_1, \zeta_1, \widehat{\eta}_1, \widehat{\zeta}_1)} . \quad (\text{A.11})$$

So finally, we have obtained, observing that

$$\delta_{3,0} = \underline{\delta}_{3,2} \widehat{\delta}_{2,1} \bar{\delta}_{1,0} ,$$

and consequently

$$\delta_{3,0} = -|\eta_3|^{1-\ell} |\eta_1|^{\ell-1} \frac{\Delta_\ell(\eta_3, \zeta_3, \widehat{\eta}_2, \widehat{\zeta}_2)}{\Delta_\ell(\eta_2, \zeta_2, \widehat{\eta}_2, \widehat{\zeta}_2)} \frac{\Delta_\ell(\eta_2, \zeta_2, \widehat{\eta}_1, \widehat{\zeta}_1)}{\Delta_\ell(\eta_1, \zeta_1, \widehat{\eta}_1, \widehat{\zeta}_1)} . \quad (\text{A.12})$$

Let us recapitulate what we have obtained in the case of the three-broken line.

(i) For $\tau \in [0, \tau_1(t, s)[$, the Jacobian is one :

$$\delta^{(1)} = \delta_{1,0} = 1 . \quad (\text{A.13})$$

(ii) For $\tau \in]\tau_1(t, s), \tau_2(t, s)[$, the Jacobian (see the main text) is

$$\delta^{(2)} := \delta_{2,0} = -|\eta_2|^{1-\ell} |\eta_1|^{\ell-1} \frac{\Delta_\ell(\eta_2, \zeta_2, \widehat{\eta}_1, \widehat{\zeta}_1)}{\Delta_\ell(\eta_1, \zeta_1, \widehat{\eta}_1, \widehat{\zeta}_1)} . \quad (\text{A.14})$$

(iii) For $\tau \in]\tau_2(t, s), +\infty[$, the Jacobian is given by $\delta^{(3)} = \delta_{3,0}$.

Generalization

It is now not too difficult to extend the formula in the case of N reflexions.

$$\delta^{(N+1)} = -|\eta_{N+1}|^{1-\ell} |\eta_1|^{\ell-1} \prod_{j=1}^N \left(\frac{\Delta_\ell(\eta_{j+1}, \zeta_{j+1}, \widehat{\eta}_j, \widehat{\zeta}_j)}{\Delta_\ell(\eta_j, \zeta_j, \widehat{\eta}_j, \widehat{\zeta}_j)} \right). \quad (\text{A.15})$$

References

- [De] M. Derridj. Subelliptic estimates for some systems of complex vector fields. In “Hyperbolic problems and regularity questions”. Series Trends in Mathematics. Edtrs : M. Padula and L. Zanghirati. Birkhäuser (2006), p. 101-108.
- [HeNi] B. Helffer and F. Nier. Hypoelliptic estimates and spectral theory for Fokker-Planck operators and Witten Laplacians. Lecture Notes in Math. 1862, Springer Verlag, Berlin 2005.
- [HeNo] B. Helffer and J. Nourrigat. *Hypoellipticité maximale pour des opérateurs polynômes de champs de vecteur*. Progress in Mathematics, Birkhäuser, Vol. 58 (1985).
- [Ho1] L. Hörmander. Hypoelliptic second order differential equations. Acta Mathematica 119 (1967), p. 147-171.
- [Ho2] L. Hörmander. Subelliptic operators. Seminar on singularities of solutions of partial differential equations. Ann. Math. Studies 91 (1978), p. 127-208.
- [JoTre] J.L. Journé and J.M. Trépreau. Hypoellipticité sans sous-ellipticité : le cas des systèmes de n champs de vecteurs complexes en $(n + 1)$ -variables. Séminaire EDP in Ecole Polytechnique, April 2006.
- [Ko1] J. Kohn. Lectures on degenerate elliptic problems. Pseudodifferential operators with applications, C.I.M.E., Bressanone 1977, p. 89-151 (1978).
- [Ko2] J.J. Kohn. Hypoellipticity and loss of derivatives, with an appendix by M. Derridj and D. Tartakoff. Ann. of Math. 162(2), p. 943-986 (2005).
- [Mai1] H.M. Maire. Hypoelliptic overdetermined systems of partial differential equations. Comm. Partial Differential Equations 5 (4), p. 331-380 (1980).

- [Mai2] H.M. Maire. Résolubilité et hypoellipticité de systèmes surdéterminés. Séminaire Goulaouic-Schwartz 1979-1980, Exp. V, Ecole Polytechnique (1980).
- [Mai3] H.M. Maire. Necessary and sufficient condition for maximal hypoellipticity of $\bar{\partial}_b$. Unpublished (1979).
- [Mai4] H.M. Maire. Régularité optimale des solutions de systèmes différentiels et du Laplacien associé : application au $\square_b$. Math. Ann. 258, p. 55-63 (1981).
- [No1] J. Nourrigat. *Subelliptic estimates for systems of pseudo-differential operators*. Course in Recife (1982). University of Recife.
- [No2] J. Nourrigat. Systèmes sous-elliptiques. Séminaire Equations aux Dérivées Partielles, 1986-1987, exposé V, Ecole Polytechnique (1987).
- [No3] J. Nourrigat. Subelliptic systems II. Inv. Math. 104 (2) (1991), p. 377-400.
- [Tr1] F. Trèves. A new method of proof of the subelliptic estimates. Comm. Pure Appl. Math. 24 (1971), p. 71-115.
- [Tr2] F. Trèves. Study of a model in the theory of complexes of pseudo-differential operators. Ann. of Math. (2) 104, p. 269-324 (1976). See also erratum: Ann. of Math. (2) 113, p. 423 (1981).